

L2 ANSWER 12 OF 27 BIOSIS COPYRIGHT (c) 2010 The Thomson Corporation on
STN

AN 2001:73648 BIOSIS

DN PREV200100073648

TI Treatment with glucocorticoids decreases both Abetax-40 and Abetax-42 in
cerebrospinal fluids.

AU Tokuda, T. [Reprint author]; Oide, T.; Tamaoka, A.; Matsuno, S.;
Hashimoto, T.; Shoji, S.; Ikeda, S.

CS Shinshu Univ Sch Med, Matsumoto, Japan

SO Society for Neuroscience Abstracts, (2000) Vol. 26, No. 1-2, pp.
Abstract No.-82.1. print.
Meeting Info.: 30th Annual Meeting of the Society of Neuroscience. New
Orleans, LA, USA. November 04-09, 2000. Society for Neuroscience.
ISSN: 0190-5295.

DT Conference; (Meeting)
Conference; Abstract; (Meeting Abstract)

LA English

ED Entered STN: 7 Feb 2001
Last Updated on STN: 12 Feb 2002

SO Society for Neuroscience Abstracts, (2000) Vol. 26, No. 1-2, pp.
Abstract No.-82.1. print.
Meeting Info.: 30th Annual Meeting of the Society of Neuroscience. New
Orleans, . . .

AB Epidemiologic studies have shown that anti-inflammatory medications
decrease the incidence of Alzheimer's disease (AD).
Corticosteroids are one of the most effective anti-inflammatory drugs.
Here we examined the concentrations of amyloid beta-protein (Abeta)
species in the cerebrospinal fluid (CSF) in the patients who were treated
with prednisolone for neuroimmunological disorders. They did
not have any clinical signs of dementia. The daily dose of
prednisolone was more than 30 mg as starting dosage, then
gradually tapered. We sequentially measured concentrations of Abeta
species in the CSF using ELISA system. The concentrations of both CSF
Abetax-40 and Abetax-42 decreased significantly after starting
prednisolone. In some patients, concentrations of them increased
when the dose of prednisolone was tapered. The concentration of
Abetax-40 and Abetax-42 changed in a parallel way in each patient. It is
likely that our patients without dementia have a normal
clearance pathway of Abeta species from the brain to CSF. We can
therefore presume that the concentrations. . . reflect the production
rate of these Abetas in the brain. Together with this, our results
suggested that moderate or high-dose prednisolone treatment
could decrease intracerebral production of Abeta species that might be
useful for the treatment of AD.